



1 21 May 2015  
2 EMA/CHMP/338679/2014  
3 Committee for Human Medicinal Products (CHMP)

4 Questions and answers on sodium in the context of the  
5 revision of the guideline on 'Excipients in the label and  
6 package leaflet of medicinal products for human use'  
7 (CPMP/463/00 Rev.1)  
8 Draft

9

Draft agreed by Excipients Drafting group	11 May 2015
Adopted by CHMP for release for consultation	21 May 2015
Start of public consultation	24 June 2015
End of consultation (deadline for comments)	30 September 2015
Agreed by Excipients Drafting group	<Month YYYY>
Adopted by CHMP	<DD Month YYYY>

10

11

Comments should be provided using this [template](#). The completed comments form should be sent to [excipients@ema.europa.eu](mailto:excipients@ema.europa.eu)

12

<b>Keywords</b>	<i>Excipients, Package leaflet, Sodium, Table salt</i>
-----------------	--

13



14 Questions and answers on sodium in the context of the  
15 revision of the guideline on 'Excipients in the label and  
16 package leaflet of medicinal products for human use'  
17 (CPMP/463/00 Rev. 1)

18 **1. Background**

19 Following the European Commission decision to revise the Annex of the guideline on 'Excipients in the  
20 label and package leaflet of medicinal products for human use' (CPMP/463/00 Rev. 1) [1], a  
21 multidisciplinary group of experts involving SWP (lead), QWP, PDCO, PRAC (ex PVWP), CMD(h), VWP,  
22 BWP and BPWP was created in 2011.

23 The objective of this group is to update the labelling of selected excipients listed in the Annex of the  
24 above mentioned EC guideline, as well as to add new excipients to the list, based on a review of their  
25 safety. The main safety aspects to be addressed were summarised in a concept paper published in  
26 March 2012 [2].

27 Draft questions and answers (Q&A) documents on excipients are progressively released for public  
28 consultation. They include proposals for new or updated information for the label and package leaflet.

29 When one or several Q&As have been finalised, the new information in the package leaflet will be  
30 included in a revised annex of the guideline.

31 For more information see the [Excipients labelling webpage](#) on the EMA website.

32 **2. What is sodium and why is it used as an excipient?**

33 Sodium is an essential nutrient. It is the principal cation in the extracellular fluid (ECF) and has a key  
34 role in maintaining fluid balance, acid base balance, muscle and nerve activity and transport of  
35 nutrients across cell membranes. The main source of dietary sodium is sodium chloride (which the  
36 majority of people know as common table salt).

37 Sodium is used in medicines both as the main ingredient (for example where medicines are intended to  
38 replace physiological sodium, or the active pharmaceutical ingredient is a sodium salt), and also as an  
39 excipient. The former is out of scope of this Q&A.

40 Sodium salts used as excipients are most commonly used to increase solubility, but they can also be  
41 used for disintegration, chelation, lubrication, binding, emulsifying and stabilising. They may be also  
42 added for colouring or for antimicrobial properties.

43 **3. Which medicinal products contain sodium?**

44 Sodium can be found in effervescent medicines. In those medicines, sodium-containing salts, including  
45 sodium bicarbonate and sodium carbonate, are commonly used with acidic agents such as citric or  
46 tartaric acid, to cause a reaction in water that produces carbon dioxide (CO<sub>2</sub>). The CO<sub>2</sub> leads to the  
47 resultant fizzing of the effervescent powder [3]. Large quantities of sodium salts may be required to  
48 enhance the solubility of a medicine. Most medicines that contain high levels of sodium are therefore

49 likely to be effervescent or soluble, however, there may be other medicines that also contain high  
50 amounts of sodium. All sodium salts are soluble and therefore even sodium bound in complex  
51 molecules would be expected to dissolve and exert a physiological effect.

52 UK medicines information (UKMI) has prepared a list of the sodium content of a variety of medicinal  
53 products available in the UK [4]. The list is not exhaustive and includes products containing a sodium  
54 salt as an excipient or as an active ingredient. The list is useful for illustrating that the sodium content  
55 of medicines is variable and also raises the following issues.

56 1) The sodium content of medicines can be very high. For example the maximum daily dose of an  
57 effervescent medicine can contain 175 mmol of sodium (201% of the WHO recommended  
58 maximum daily intake for sodium for an adult).

59 2) Many medications that are high in sodium are commonly used for a wide variety of conditions  
60 and are often available over the counter (OTC) without a prescription. This reduces the  
61 opportunity for pharmacists and other health care professionals to advise people on sodium  
62 levels.

63 3) People who have a preference or need for medicines that dissolve, may be taking several  
64 medicinal products in a dissolvable form. This additive effect could have a significant impact on  
65 their sodium intake.

#### 66 **4. What are the safety concerns?**

67 Increasing the level of sodium in the body causes an expansion of ECF which increases blood pressure  
68 (BP) [5]. Maintaining steady sodium levels is principally achieved through regulation of excretion  
69 through the kidneys (renal excretion) [5, 6]. The capacity for renal excretion is lower in the very young  
70 and the elderly [5].

71 High intake of table salt is associated with high BP (hypertension) and stroke in adults [5, 6].

72 A study in 1,292,337 patients over the age of 18 recently reported that the high sodium content of  
73 some effervescent, soluble and dispersible medicines might be associated with an increased risk of  
74 cardiovascular disease [7]. Risk was measured using a statistic known as the odds ratio and presented  
75 including a confidence interval which shows the range of risks that may be true. In the study, patients  
76 who survived either a heart attack or a stroke or who died of a cardiovascular condition were 1.16  
77 (1.12–1.21) times more likely to have been prescribed an effervescent, soluble or dispersible medicine  
78 than patients who did not suffer one of these events. The patients who survived a stroke were 1.22  
79 (1.16–1.29) times more likely to have been prescribed a medicine that dissolves and patients with high  
80 BP (hypertension) were 7.18 (6.74–7.65) times more likely to have had prescriptions for dissolvable  
81 medicines. All of these results were statistically significant indicating that there is likely to be a true  
82 association between the high sodium content of effervescent, dispersible and soluble medicines and  
83 cardiovascular disease.

84 The study has been criticised because it did not look separately at patients who took medicines for a  
85 long time and those who only took medicines for a very short length of time. Those patients taking  
86 medicines regularly and for a prolonged period of time might be expected to be at higher risk from  
87 sodium in medicines than those patients taking only a short-term course of treatment. Another  
88 criticism of the study is that it was not able to consider the amount of sodium patients were having in  
89 their diets as this information is not available in the database that was used for the study. There are  
90 likely to have been differences in the amounts of sodium patients were having in their diet which,  
91 independent of the salt in their medicines, may have affected their risk of cardiovascular disease.

92 Despite the criticisms of the study, the association between hypertension and high sodium-containing  
93 effervescent, soluble and dispersible medicines was very strong, and so is likely to represent a true  
94 association. This is supported by the fact that an association is already accepted to exist between high  
95 dietary sodium and cardiovascular events, especially hypertension.

96 Increasing long-term intake of dietary sodium has been shown to increase BP across all study  
97 populations and age ranges [8]. Prolonged high BP has been associated with stroke, myocardial  
98 infarction, heart failure and kidney disease and has also been linked to dementia and premature  
99 death [9]. Several large studies (meta-analyses) have reported an increased risk of stroke with  
100 increased dietary sodium [10–13]. Reducing salt has been shown to significantly reduce BP [12, 14–  
101 16] and lower BP has been proven to reduce cardiovascular disease. The effect of low salt diets on BP  
102 is not always maintained beyond 6 months, but this is felt to reflect the difficulty in maintaining a low  
103 salt diet, rather than salt reduction having only a short-term effect on BP [5].

104 Young babies have lower capacity for removing sodium from the body. Acute high sodium intakes from  
105 any source can result in dangerously raised sodium levels in the blood (hypernatraemia). This can  
106 result most commonly in events including listlessness, serious dehydration and seizures [17].

107 Chronic high dietary sodium intake can raise blood pressure in children which increases the risk of  
108 hypertension and cardiovascular disease in adulthood [5, 6]. High dietary sodium intakes in childhood  
109 are also associated with the development of a preference in later life for salty food [18].

110 The WHO recommend adults to consume less than 5g of sodium chloride (table salt) per day  
111 (equivalent to less than 2g (or 87 mmol) sodium per day). Individual countries have their own  
112 guidance; for example in the UK it is recommended for adults to have less than 6g of sodium chloride  
113 per day (equivalent to less than 2.4g (or 104mmol) sodium per day).

114 For children, the WHO advise that recommended maximum daily intakes should be proportional to  
115 adults and based on energy requirements [6]. An EU Framework on Voluntary National Salt  
116 initiatives [19] has been agreed at population level, in order to achieve the national or WHO  
117 recommendations.

## 118 **5. What are the reasons for updating the information in the** 119 **package leaflet?**

120 It is important that patients, parents, carers, pharmacists, prescribers and other healthcare  
121 professionals are able to easily identify how much sodium is present in medicines, especially given that  
122 many medicines affected by this issue are available OTC with minimal chance for a healthcare  
123 professional to offer any advice on sodium and potential risks.

124 Information on sodium should be readily available to everyone in an understandable format in the  
125 product information and will allow more informed decisions to be made about whether a medicine and  
126 its ingredients are appropriate for the patient. This information will be particularly important for people  
127 who are on sodium restricted diets or have pre-existing cardiovascular disease and especially those  
128 who need to take medicines on a regular basis.

129 The concerns with the current labelling of sodium as an excipient are:

- 130 • Sodium may not be familiar to patients and parents as being part of sodium chloride and the main  
131 component of dietary salt (common table salt).
- 132 • The threshold level below which a medicine is considered sodium-free is 1mmol / dose. This is a  
133 low level of sodium and only 1.1% of the WHO recommended maximum daily intake. Any

134 medicine with more than 1 mmol of sodium per dose includes a warning that its use should be  
135 taken into consideration by patients on a low salt diet. No distinction is made between medicines  
136 that contain relatively low levels of sodium and those that contain exceedingly high levels of  
137 sodium (e.g. approx. 22 mmol / dose for some effervescent products meaning each dose  
138 represents 25% of the WHO recommended maximum daily amount for sodium). It is therefore  
139 difficult for patients and prescribers to appreciate which medicines have particularly high levels of  
140 sodium.

- 141 • Sodium is presented in units of mmol or mg and neither may be meaningful to patients or  
142 prescribers.

143 The high levels of sodium in some medicines may not be appreciated by patients or healthcare  
144 professionals.

145 The proposed updates to the labelling of sodium therefore include defining and presenting sodium  
146 content in a clear and meaningful way. Thus the sodium content in the maximum daily dose  
147 recommended for a medicine will be presented as a proportion of the WHO maximum recommended  
148 daily dietary intake for sodium. So for example, if one tablet contains 250mg (or about 11mmol)  
149 sodium and a maximum of four tablets may be taken in a day, this corresponds to a maximum daily  
150 dose of 1g of sodium (approximately 44mmol). This would be the equivalent to approximately 50% of  
151 the 2g (or 87mmol) of sodium that the WHO recommends to be the maximum daily dietary intake for  
152 an adult. A patient or healthcare professional will be able to clearly see that the maximum daily dose of  
153 the medicine provides half of their maximum daily recommended intake of sodium.

154 The proposal also introduces an additional threshold to define levels of sodium in medicines considered  
155 to be 'high'. There is no evidence to suggest what level of sodium in medicines is acceptable and this  
156 will vary by individual. However, it is proposed that any product where the maximum daily dose  
157 contains  $\geq 17$  mmol (391 mg) sodium (approximately 20% of the WHO recommended maximum daily  
158 intake for sodium), should be considered as having a 'high' sodium content. This is an empirical figure  
159 but is based on the fact that the intake of sodium through medicines is in addition to dietary sodium  
160 and many people are already consuming too much sodium through common salt in their diet.

161

162 **Information in the package leaflet as per the 2003 Guideline**

Name	Route of Administration	Threshold	Information for the Package Leaflet	Comments
Sodium	Parenteral	Less than 1 mmol per <dose>	This medicinal product contains less than 1 mmol sodium (23 mg) per <dose>, i.e. essentially 'sodium- free'.	Information relates to a threshold based on the total amount of Na <sup>+</sup> in the medicinal product. It is especially relevant to products used in paediatric doses, to provide information to prescribers and reassurance to parents concerning the low level of Na <sup>+</sup> in the product.
	Oral Parenteral	1 mmol per <dose>	This medicinal product contains x mmol (or y mg) sodium per dose. To be taken into consideration by patients on a controlled sodium diet	

163 **5. Proposal for an updated information in the package leaflet**

Name	Route of Administration	Threshold*	Information for the Package Leaflet	Comments
<b>Sodium</b>	Parenteral	Less than 1 mmol per dose	This medicinal product contains less than 1 mmol sodium (23 mg) per <dose>, i.e. it is essentially 'sodium- free'.	1 mmol of sodium (Na) = 23 mg Na = 57 mg table salt (NaCl).
	Oral Parenteral	1 mmol (23 mg) per dose	One dose of this medicinal product contains <X mg> sodium (found in table salt). This is equivalent to <Y%> of the recommended maximum daily intake of sodium for an adult.	
	Oral Parenteral	17 mmol (391 mg) in the maximum daily dose	The maximum recommended daily dose of this medicinal product contains <X mg> sodium (found in table salt). This is equivalent to <Y%> of the recommended maximum daily intake of sodium for an adult.  Talk to your pharmacist or doctor if <you> <or> <your child> need(s) [product name] on a daily basis for a prolonged period of time, especially if <you><they> have been advised to follow a low salt diet.	17mmol (391mg) is approximately 20% of the WHO adult recommended maximum daily dietary intake for sodium and is considered to represent 'high' sodium.  This is also relevant for children, where the maximum daily intake is considered to be proportional to adults and based on energy requirements.

164  
165 \* Note: The threshold is a value, equal to or above which it is necessary to provide the information stated [1].

## 166 References

- 167 1. Guideline on excipients in the label and package leaflet of medicinal products for human use'  
168 (CPMP/463/00 Rev.1). July 2003.  
169 [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2009/09/WC50](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC50)  
170 [0003412.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2009/09/WC500003412.pdf)
- 171 2. Concept paper on the need for revision of the 'Guideline on excipients in the label and package  
172 leaflet of medicinal products for human use' (CPMP/463/00) EMA/CHMP/SWP/888239/2011  
173 [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Scientific\\_guideline/2012/03/WC50](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/03/WC50)  
174 [0123804.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Scientific_guideline/2012/03/WC500123804.pdf)
- 175 3. Martindale. The complete drug reference. 37th edition. Pharmaceutical Press.
- 176 4. UKMi. Medicines Q&As. Q&A 145.4; What is the sodium content of medicines? 2012  
177 <https://www.evidence.nhs.uk/search?q=%22What+is+the+sodium+content+of+medicines%22>  
178 [2](https://www.evidence.nhs.uk/search?q=%22What+is+the+sodium+content+of+medicines%22)
- 179 5. Salt and Health Scientific Advisory Committee on Nutrition. 2003
- 180 6. World Health Organisation 2012. Guideline: Sodium intake for adults and children.
- 181 7. George J, Majeed W, Mackenzie I, MacDonald T, Wei L. The Association of Cardiovascular  
182 Events with Sodium-Containing Effervescent, Dispersible and Soluble Medications; Nested  
183 Case-control Study. BMJ 2013.
- 184 8. Intersalt Cooperative Research Group. Intersalt: an international study of electrolyte excretion  
185 and blood pressure. Results for 24 hour urinary sodium and potassium excretion. BMJ  
186 1988;297:319-28
- 187 9. Hypertension: Clinical management of primary hypertension in adults. NICE guideline CG127  
188 <http://www.nice.org.uk/guidance/cg127>
- 189 10. Li XY, Cai XL, Bian PD, Hu LR. High salt intake and stroke: meta-analysis of the epidemiologic  
190 evidence. CNS Neurosci Ther 2012;18(8):691-701.
- 191 11. O'Donnell Martine et al. Urinary Sodium and Potassium Excretion, Mortality, and Cardiovascular  
192 Events N Engl J Med 2014;371:612-623.  
193
- 194 12. Aburto NJ, Ziolkovska A, Hooper L, Elliott P, Cappuccio FP, Meerpohl JJ. Effect of lower sodium  
195 intake on health: systematic review and meta-analyses. BMJ 2013;346:f1326.
- 196 13. Strazullo P, D'Eila L, Kandala NB et al. Salt intake, stroke and cardiovascular disease: meta-  
197 analysis of prospective studies. BMJ, 2009, 339:b4567.
- 198 14. MacGregor GA, He FJ. Effect of modest salt reduction on blood pressure: a meta-analysis of  
199 randomized trials. Implications for public health. Journal of Human Hypertension 2002;16:761-  
200 770
- 201 15. He FJ, MacGregor GA. Effect of longer-term modest salt reduction on blood pressure. Cochrane  
202 Database of Systematic Reviews, 2013,(4):CD004937
- 203 16. Dickinson HO, Mason JM, Nicolson DJ et al. Lifestyle interventions to reduce raised blood  
204 pressure: a systematic review of randomized controlled trials. J Hypertens, 2006, 24(2):215-  
205 233



- 206 17. Hypernatraemia. Professional reference article. Patient.co.uk 20<sup>th</sup> February 2012  
207 <http://www.patient.co.uk/doctor/Hypernatremia.htm>
- 208 18. World Action On Salt and Health How does salt affect children?  
209 <http://www.worldactiononsalt.com/salthealth/children/>
- 210 19. European Union framework for national salt initiatives. European Commission. 2008.